



Food Handler Basic Course Study Guide



Table of Contents

Why Read This Manual?	1	4.5 Parasites	16
Food Handler Card.....	1	4.6 Chemicals.....	16
Training Required	1	4.7 Physical Contamination.....	16
Reciprocity and Equivalency.....	1	4.8 Contaminated Food.....	16
How to Use This Manual.....	1		
A “Person in Charge” is Required.....	1		
Food Safety Manual.	2	Chapter 5. Food Temperatures.....	17
Chapter 1. Food Handler Training		5.1 Temperature Control (for Food Safety) ..	17
Learning Objectives	3	5.2 The “Danger Zone”.....	17
1.1 Employee Illness.....	3	5.3 When to Discard Food.....	17
1.2 Hand-Washing.....	3	5.4 Cooking Food.....	17
1.3 No Bare Hand Contact with		5.5 When is Cooked Food Safe?	18
Ready to Eat Food	4	5.6 Remember	18
1.4 Your Role in Helping Prevent		5.7 Cold Holding.....	18
Foodborne Illness	5	5.8 Hot Holding.....	18
1.5 The Role of Management in Helping		5.9 Keep it Hot	18
Prevent Foodborne Illness.....	6	5.10 Keep it Cold	18
1.6 Foodborne Illness	6	Chapter 5 Review	19
1.7 Temperature Control	7	Chapter 6. Safe Storage Practices	20
1.8 Final Cooking Temperature.....	7	6.1 Cross Contamination	20
1.9 Contamination and		6.2 Keep Foods Safe from Contamination ...	20
Cross Contamination	8		
1.10 Work Only When You Are Well	9	Chapter 7. A Clean Workplace is Safer.....	21
Chapter 1 Review	10	7.1 Follow These Important Rules.....	21
Chapter 2. Prevent the Spread of Disease.....	11	7.2 Utensils, Surfaces and Equipment.....	21
2.1 Why Hand Washing is Very Important.....	11	7.3 Directions.....	22
2.2 Germs are Everywhere	11	7.4 Remember	22
2.3 No Bare Hand Contact.....	12	Chapter 7 Review	23
2.4 Ready-To-Eat.....	12		
2.5 Gloves and Hand-Washing	12	Glossary	24
Chapter 2 Review	13		
Chapter 3. Employee Practices.....	14	Reference.....	25
3.1 Take Care of How You Look		General.....	25
and How You Act	14	Employee Illness.....	26
3.2 Fingernails	14	Most Common Illnesses from Employees	26
3.3 Gloves Can Spread Germs	14	Hand-Washing	27
3.4 Beverages	14	Cooking Temperatures	28
		“Is It Done Yet?”	30
Chapter 4. What Makes People Sick		Calibrating a Dial Food Thermometer.....	32
from Food?.....	15	Cooling Hot Foods.....	33
4.1 Foodborne Illness	15	Date Marking	36
4.2 Potentially Hazardous Foods		Pests	36
(Time/Temperature Control).....	15		
4.3 Bacteria.....	15		
4.4 Viruses.....	16		

Why Read This Manual?

Food Handler Card

All food handlers employed in food service must obtain a food handler card within 30 days from the date of hire. As a food handler, you are required to keep your food handler card current by renewing it every three years or as required.

Training Required

The goal of this program is to provide you with a basic understanding of food safety. This will assist your manager, who is responsible for ensuring that you prepare and serve food safely. **A food handler card confirms that you have met the learning objectives for this certificate course.**

Reciprocity and Equivalency

Any person who has a current certificate from a department-approved food manager certification program need not obtain a food handler card.

To be accepted in place of a food handler card, a food manager certificate must be current and renewed upon expiration.

Photocopies of the food handler cards and food manager training certificates should be kept at the facility to show the health inspector upon request.

How to Use This Manual

This manual is intended to help you learn what you need to know to obtain a food handler card. You will need a score of 70% to pass. You will be tested on all of the learning objectives that are listed on pages (3-9). Throughout this manual you will find study questions that will help you get ready to take the test for the food handler card.

The words that are ***italicized and bold*** are explained in the glossary located in the back of this manual.

A “Person in Charge” is Required

Someone at your restaurant must be in charge during all hours of operation. This person in charge (PIC) is responsible for knowing the food sanitation rules and the procedures within your establishment. This person is responsible for providing you with information you need to perform your job.

The PIC is usually a manager or supervisor, but can be anyone who can demonstrate the knowledge listed above, and is given the authority to oversee other employees.



Keep this manual as a
reference and download the
Food Safety Manual.

Chapter 1. Food Handler Training Learning Objectives

Food workers are expected to know this information to obtain their food handler card.

The concept of foodborne illness will be introduced. The training will address personal hygiene, contamination, and temperature control to reinforce the food handler's behaviors, which can prevent foodborne illness.

Section 1.1 Employee Illness

- The food handler will know to call the person in charge at the food service facility when ill with diarrhea, vomiting, jaundice, or fever with sore throat.
- The food handler will know not to work in the food service facility while ill with these symptoms.
- The food handler will know to not work in food service for 24 hours after symptoms of diarrhea or vomiting have gone.
- The food handler will know not to handle food with an infected boil, cut, burn, or sore on the hand or wrist. Food may be handled if the injury is covered with a clean bandage and a latex-free glove.

Section 1.2 Hand-Washing

Workers will understand elements of good hand-washing.

- The food handler will be able to identify the correct technique for hand-washing:
 - Use running warm water and soap
 - Scrub hands and rinse thoroughly (approximately 20 seconds)
 - Dry hands with single-use towel, or air dryer



- The food handler will be able to identify situations when food handlers must wash their hands:
 - Before starting work
 - After using the toilet and again when entering work area
 - After handling raw food and raw animal products
 - After handling dirty dishes
 - After handling garbage
 - After cleaning or using chemicals
 - After blowing nose, sneezing, coughing, or touching eyes, nose or mouth
 - After smoking, or using tobacco products
 - After eating or drinking
 - Before putting on food service gloves
- The employee food handler will know that the food service gloves can spread germs and is not a substitute for proper hand washing.
- The food handler will know that smoking, eating, and chewing tobacco is prohibited in food preparation areas, including food and utensil storage areas.

Section 1.3 No Bare Hand Contact with Ready to Eat Food

- The food handler will know that touching ready-to-eat foods with bare hands is not allowed. A food worker must use utensils or single use gloves when working with ready-to-eat food. Good hand washing is still required even when using utensils or gloves to handle food.
- The food handler will know that food service gloves are capable of spreading germs and are not a substitute for proper hand-washing.



- The food handler will be able to identify situations when gloves must be discarded; hands washed and new gloves are worn:
 - As soon as gloves become soiled or torn
 - Before beginning a different task
 - After handling raw meat, fish, or poultry
 - Before handling ready-to-eat food.
- The food handler will be able to identify ready-to-eat food as foods that are edible without washing, cooking, or additional preparation. These foods include:
 - Raw, washed and cut fruits and vegetables; and
 - Food that require no additional cooking
- The food handler will know they may prevent bare-hand contact by using:
 - Deli tissue
 - Spatulas/Tongs/Forks/Utensils
 - Dispensing equipment
 - Single-use gloves

Section 1.4 Your Role in Helping Prevent Foodborne Illness

- The food handler will be able to describe five major mistakes that often cause foodborne illness:
 1. Inadequate hand-washing
 2. Employees working while they are ill
 3. Cross contamination
 4. Inadequate cooking temperatures
 5. Inadequate temperature control
(allowing foods to be in the danger zone)



- The food handler will be able to describe the activities performed by food handlers that prevent foodborne illness from happening. Activities preventing foodborne illness include:
 - Proper hand-washing every time hands may have become contaminated
 - Food handlers working only when healthy
 - Storing and handling of foods in a manner to prevent contamination
 - Cooking each animal product to its required internal temperature
 - Maintaining hot and cold temperatures (keeping foods out of the danger zone)
 - Use of utensils or single-use gloves to prevent bare hand contact with ready-to-eat food.

Section 1.5 The Role of Management in Helping Prevent Foodborne Illness

- The food handler will know that the manager sets the tone of what food safety activities occur or don't occur within the facility.
- The food handler will know that the food service management is responsible for training and ensuring that food handlers practice activities that prevent foodborne illness.

Section 1.6 Foodborne Illness

- The food handler will be able to describe foodborne illness as an illness resulting from eating contaminated food.
- The food handler will know that food contaminated with organisms (germs) does not always look, smell or taste different from non-contaminated food.
- The food handler will know that symptoms vary and may include diarrhea, vomiting, fever, cramping and nausea.
- The food handler will know that depending on the cause, symptoms may develop in a few minutes to several days. Some symptoms may last several days and can result in death.
- The food handler will know that foodborne illness is caused by organisms (germs), chemicals, or toxins.

Section 1.7 Temperature Control

Workers will understand why hot and cold holding temperatures are important factors in preventing illness.

- The food handler will be able to identify potentially hazardous foods as food that will support bacterial growth when held at temperatures in the danger zone.
- The food handler will be able to identify the danger zone as any temperature between 41°F and 135°F.
- The food handler will be able to identify that food being cooled or heated must move through the danger zone as rapidly as possible.
- The food handler will be able to identify 135°F or hotter as the proper temperature for hot holding potentially hazardous food.
- The food handler will be able to identify 41°F or colder as the proper temperature for cold holding potentially hazardous food.
- The food handler will know that you cannot make food safe to eat when food has been in the danger zone for four hours or more.

Section 1.8 Final Cooking Temperature

Workers will understand why cooking foods to proper temperatures are important for preventing illness.

The food handler will be able to identify that cooking foods to the recommended temperature will kill disease-causing germs.



Section 1.9 Contamination and Cross Contamination

Workers will understand why cross contamination is dangerous and know ways to prevent it.

- The food handler will be able to define and identify physical contamination as foreign objects accidentally introduced into food. Food items may arrive already contaminated with dirt and pebbles.
- The food handler will be able to define and identify cross contamination as happening when microorganisms are transferred from one food or surface to another food.
- The food handler will be able to identify methods to prevent cross contamination such as washing, rinsing, and sanitizing utensils, work surfaces and equipment between uses.
- The food handler will be able to identify storage conditions that will minimize the potential for cross contamination:
 - Store raw meats below and completely separate from ready-to-eat food in refrigeration units
 - Store chemicals, cleansers and pesticides completely separate from food, utensils, and single service items
 - Properly label all chemicals, cleansers and pesticides



Section 1.10 Work Only When You Are Well

If you feel sick you should not go to work. The germs you bring to work can spread when you touch food, dishes, counters, utensils, and other people.

- Do not work if you have a fever and sore throat
- Do not work if you have loose bowels (diarrhea)
- Do not work if you are throwing up (vomiting)
- Do not work if you have yellowing of the skin or dark tea colored urine (jaundice)
- Wait for at least 24 hours after vomiting or diarrhea have stopped before returning to work.



Tell your manager if you have any of these symptoms. If the manager has questions, he or she can call the County Health Department.

Do not handle food with an **infected** boil, burn, cut or sore on your hand. Food may be handled if you cover the injury with a clean bandage, and wear a latex-free glove.

Chapter 1 Review

At the end of each chapter, we've included some review questions to help you make sure you understand the material. We strongly recommend that you complete them as you read through the manual.

1. When you have a sore throat with fever or diarrhea, you should:
 - A. Go to work and tell your coworkers to be careful around you
 - B. Call your manager and report that you are sick
 - C. Take medicine to stop the symptoms and go to work
 - D. Not tell anyone and continue working
-

2. If you are ill, tell your manager before starting work.
 - A. True
 - B. False
-

3. Food handlers can contaminate food when they:
 - A. Discard food left in the danger zone
 - B. Take a break in a designated area
 - C. Keep chemicals away from food
 - D. Have infected wounds or injuries
-

4. Food handlers can contaminate food when they:
 - A. Keep fingernails short for easy cleaning underneath
 - B. Change gloves for fresh gloves between tasks
 - C. Have been diagnosed with a foodborne illness
 - D. Drink from a closed beverage cup with lid and straw

Chapter 2. Prevent the Spread of Disease

Section 2.1 Why Hand Washing is Very Important

Wash your hands often when working with food and drinks – this gets rid of germs that can make people sick. Wash your hands for approximately 20 seconds with warm running water and soap, and then dry them with clean paper towels, or an air dryer.



Remember to always wash your hands:

- **Before** you start preparing food
- **After** using the toilet and again when entering work area
- **After** handling raw animal products
- **After** handling dirty dishes and garbage
- **After** cleaning or using chemicals
- **After** blowing nose, sneezing, coughing, or touching eyes, nose or mouth
- **After** smoking or using tobacco products
- **After** eating or drinking
- **Before** you put on latex-free gloves

Section 2.2 Germs are Everywhere

Germs such as **bacteria** and **viruses** are everywhere. If you do not wash your hands in the right way and use a barrier between your hands and ready-to-eat food, your hands can put germs in food that will be eaten by your customers. Your customers may get sick from these germs.

Section 2.3 No Bare Hand Contact

Bare hand contact with ready-to-eat food is not allowed. Ready-to-eat foods are edible without washing, cooking or additional preparation before service.

Section 2.4 Ready-To-Eat

These foods include:

- Raw, washed and cut fruits and vegetables
- Food that require no additional cooking, such as sandwiches and salads

Instead of bare hands, you can use:

- Utensils, tongs, forks
- Deli tissue
- Dispensing equipment or
- Non-latex, single-use gloves

Section 2.5 Gloves and Hand-Washing

Gloves and other barriers do not replace hand-washing. Wash hands before putting gloves on and when changing to a new pair.

Change your gloves:

- As soon as they become soiled or torn
- Before beginning a different task
- After handling raw meats, fish, or poultry and
- Before handling ready-to-eat food

Chapter 2 Review

1. You must wash your hands between changing gloves?
 - A. True
 - B. False

2. When must you wash your hands?
 - A. After sneezing or coughing
 - B. After touching raw meat
 - C. After eating or drinking
 - D. All of the above

3. No bare hand contact is required when foods requiring no additional cooking are served.
 - A. True
 - B. False

4. When must you change single-use gloves?
 - A. Every hour
 - B. At the beginning of the shift
 - C. Before handling ready-to-eat food
 - D. Before handling raw meats

Chapter 3. Employee Practices

Section 3.1 Take Care of How You Look and How You Act

Do not smoke or chew tobacco while you are working or when you are near food or dishwashing areas. Smoke only while you are on a break. After you smoke, wash your hands before you return to work.

Section 3.2 Fingernails

Be sure to scrub underneath your fingernails. It is much easier to keep fingernails clean when they are kept short.

Section 3.3 Gloves Can Spread Germs

Single-use food service gloves can also spread germs. Always wash and dry your hands before putting on gloves. Wash again when gloves are removed. Change gloves between tasks. When you wear gloves be aware that gloves can spread germs onto food that will not be cooked. Even when you wear gloves, it is best to keep fingernails short.

Section 3.4 Beverages

When you are thirsty while working, you may drink from a closed beverage cup with lid and straw or cup with lid and handle. This is allowed only if the food worker is careful to prevent contamination of hands, equipment, any service items, and exposed food.



Chapter 4. What Makes People Sick from Food?

Section 4.1 Foodborne Illness

People can get sick when the food they eat has germs. Germs cause foodborne illness or food poisoning.

Section 4.2 Potentially Hazardous Foods (Time/Temperature Control)

Germs grow easily in foods like meat, fish, poultry, milk, re-fried beans, cooked rice, baked potatoes and cooked vegetables. These are called potentially hazardous foods. These are all foods that are moist and they have nutrients that the germs need to grow. Germs grow well on these foods at warm temperatures between 41°F and 135°F.

Section 4.3 Bacteria

Different kinds of germs can make people sick. Bacteria are one kind of germ. They grow fast and they may cause foodborne illness. Some bacteria make toxins that act like a poison. Cooking does not destroy most toxins. Almost always, the food looks and smells good, but it may have enough bacteria or toxin to make someone sick. Toxins can occur in many foods that have not been kept cold enough or hot enough.



Section 4.4 Viruses

A virus is another kind of germ that causes illness when it gets into the food.

You can have a virus and not know it. Even before you start feeling sick, you may be passing viruses into the food by not washing your hands after coughing, sneezing or using the toilet. This is one reason why hand-washing and not touching ready-to-eat foods with your bare hands is so important. Two examples of these are the hepatitis A and norovirus.

Section 4.5 Parasites

Tiny worms that live in fish and meat are called parasites. Cooking fish and meat to the right temperature will kill parasites.

Section 4.6 Chemicals

People can also get sick when chemicals get into the food. Be sure to keep chemicals away from food.

Section 4.7 Physical Contamination

Physical contamination is when foreign objects are accidentally introduced into food. Food items may arrive already contaminated with dirt, and pebbles. Physical contamination such as broken glass can also happen at the facility.

Section 4.8 Contaminated Food

The food is contaminated. Now what? Discard contaminated food, and notify your manager right away!




Chapter 5. Food Temperatures

Section 5.1 Temperature Control (for Food Safety)

This section is about killing germs with cooking and stopping their growth by keeping the food hot or cold. This is called temperature control, and your restaurant needs at least one accurate metal-stem probe (food) thermometer to check food temperatures.

Section 5.2 The “Danger Zone”

Germs like bacteria need time, food and moisture to grow. The temperature between 41°F (5°C) and 135°F (57°C) is called the “Danger Zone!” When food sits in the “Danger Zone”, bacteria can grow fast and make toxins that can make you and others sick.



Section 5.3 When to Discard Food

Foods left in the Danger Zone for more than four hours must be discarded. Reheating the food may kill the bacteria but the toxins (produced by bacteria) will remain in the product and cause illness.

Section 5.4 Cooking Food

Cooking raw food to the proper temperature will kill germs that cause people to become sick.

Section 5.5 When is Cooked Food Safe?

Different foods have to reach different temperatures to be done or safe. Ask your manager to show you a temperature chart for cooking meats (see page 32). Be sure to cook the food to the temperature that is shown on the chart.

Section 5.6 Remember

You can choose several ways to cook food. No matter how you cook the food, it must reach the correct cooking temperature. Using a metal-stem probe thermometer is the only way to know the correct temperature of food. You must place the thermometer in the thickest part of the meat or in the center to get a true reading.

Section 5.7 Cold Holding

Always keep cold food at 41°F (5°C) or colder. Fish, shellfish, poultry, milk and red meat will stay fresh longer if you hold them cold at 41°F (5°C) or colder.



Section 5.8 Hot Holding

After the food is cooked and ready to serve, you will need to keep it warm enough to stop any germs from growing. You must turn on steam tables, soup warmers and heated surfaces before you need them so that they will be hot enough when you put the cooked food into them. Keep hot food at 135°F (57°C) or hotter.

Section 5.9 Keep it Hot

Stir food to help keep the food on top hot. A cover on the pan helps to keep the heat inside.

Section 5.10 Keep it Cold

Food being held cold on the top section of a refrigerated preparation unit also benefits from being covered.



Chapter 5 Review

1. What is it called when someone gets sick from eating food contaminated with germs or toxins?
 - A. Potentially hazardous food
 - B. Ptomaine poisoning
 - C. Foodborne illness
 - D. Raw food

2. How hot must food be kept at on the steam table to keep food safe?
 - A. Hot - 120°F
 - B. Hot - 130°F
 - C. Hot - 135°F
 - D. Hot - 165°F

3. How cold does a salad bar or refrigerator have to be to keep food safe?
 - A. Cold - 41°F
 - B. Cold - 65°F
 - C. Cold - 51°F
 - D. Cold - 55°F

Chapter 6. Safe Storage Practices

You want all the food you use to be healthy and safe. This section talks about how to safely store and handle food.

Section 6.1 Cross Contamination

Cross contamination happens when germs from raw or unclean food get into foods that are ready to serve or that will not be cooked again before you serve them.

Section 6.2 Keep Foods Safe from Contamination

As a food handler you must prevent **cross contamination**. Here are some important ways that you can prevent **cross contamination**.

- Store raw meat, fish, and poultry on the lower shelves of the refrigerator.
- Don't let raw meats; beef, pork, lamb, fish or poultry drip onto foods that will not be cooked before serving.
- Keep different types of raw meat separate from each other.
- Store unwashed food or raw food away from ready-to-eat food.
- Wash your hands between handling raw meat and foods that will not be cooked before eating.
- Never store foods that will not be cooked before serving in the same container as raw meat, fish or poultry.
- Wash your hands before working with food and before wearing gloves.
- Use utensils or disposable gloves to work with ready-to-eat food.
- Wash, rinse and sanitize the cutting surface and all the utensils and knives every time you finish with a job or between preparing different foods.
- Use clean utensils instead of hands for dispensing food.
- Store foods away from cleaners and poisons.

Chapter 7. A Clean Workplace is Safer

It takes more than soap and water to keep a food business clean and safe. You will likely be using detergents and sanitizers.

Section 7.1 Follow These Important Rules

Know what the directions say for using chemicals. Read the labels and talk to your manager about when to use them and how much to use.

Be sure you understand and follow the directions!

- Keep chemicals away from food and clean utensils. If chemicals must be stored in the same room, be sure they are stored in their own area. The area should be below food and utensils, so there is no chance of chemicals splashing onto the food and utensils.
- Can you read the labels? Are they easy to see? If they are not, ask your manager to re-label them so that everyone can read them.
- Keep all chemicals in the bottles or boxes they come in. If you put them in a different container, label them clearly.

Section 7.2 Utensils, Surfaces and Equipment

Another way to prevent cross contamination is to be sure that utensils, work surfaces and equipment are washed, rinsed, and sanitized between uses.

- Wash them in hot soapy water
- Rinse them in clean hot water
- Sanitize them with freshly prepared sanitizer (1 to 2 teaspoons of bleach per 1 gallon of water)



Section 7.3 Directions

Follow the cleaning directions for each piece of equipment.

Section 7.4 Remember

The correct steps for cleaning utensils, food contact surfaces and equipment are:

1. Wash
2. Rinse
3. Sanitize



Chapter 7 Review

1. You should store chemicals separate and away from foods?
 - A. True
 - B. False

2. Where in the refrigerator should you store raw meat?
 - A. Above the vegetables
 - B. Next to the vegetables
 - C. With the vegetables
 - D. Below the vegetables

3. The most important reason to wash, rinse and sanitize a meat slicer or knife is to:
 - A. Eliminate odors and tastes from getting into other foods
 - B. Prevent bacteria on one food from contaminating another food
 - C. Make the meat slicer or knife last longer and look better
 - D. Prevent garlic or other flavors from getting onto other foods

4. You found cooked rice left out on the counter all night, the safe way to handle it is to:
 - A. Put the rice in the refrigerator
 - B. Throw the rice into the garbage
 - C. Take a temperature of the rice
 - D. Reheat the rice to 165°F (74°C)

Glossary

Bacteria – Bacteria are germs with only one cell that can multiply into large numbers when food is in the danger zone for more than 4 hours.

Bare Hand Contact Prohibition with Ready-to-Eat Food – Bare hands may not come into contact with food that is ready to eat, such as salad or sandwiches.

Chemicals – In this manual, chemicals are referred to as ingredients in cleaning, sanitizing, or pesticide products that make people sick if eaten.

Cold Holding – Cold holding is when you keep food cold by using refrigeration or ice.

Cross Contamination – When germs from one food item are passed to another food item, usually raw food to ready-to-eat food.

Danger Zone – The Danger Zone is when the temperature of food is between 41°F (5°C) and 135°F (57°C). This is called the danger zone because bacteria will grow quickly between these temperatures.

Foodborne Illness – Sickness caused from germs or toxins in food. This is also called food poisoning.

Food Thermometer – A metal-stem probe thermometer used to take temperatures of food.

Hot holding – Holding food hot after it has been properly cooked or reheated. Food must maintain a temperature of 135°F (57°C) or hotter.

Infected – A cut or burn that is swollen, red, or has pus.

Parasites – These are tiny worms that live in fish, meat and humans.

Potentially Hazardous Foods (Time/Temperature Control for Safe Food) – Moist, nutrient-rich foods that supports the growth of bacteria when the temperature is between 41°F (5°C) and 135°F (57°C).

Reheating for Hot Holding – The process of making a cold food hot before placing on warming unit. Food must be heated from 41°F (5°C) to 165°F (74°C) within two hours.

Sanitize – The final step to removing bacteria from food contact surfaces that have just been cleaned. Many places use a solution made up of one teaspoon of bleach to one gallon of water to sanitize equipment and utensils.

Virus – Viruses are germs that can only reproduce inside of a living cell. It takes a small number of viruses to make someone sick. Many viruses get into the food from the lack of hand-washing especially after using the toilet and then touching food.

Reference

Food safety questions may come up that were not covered in the food handler training section of this manual. This reference section provides some additional food safety information.

General

Person In Charge (PIC)

The Food Sanitation Rules require that the licensee designate a person in charge (PIC) during all hours of the operation. The PIC must know foodborne illness prevention and the requirements of the Food Sanitation Rules. Every PIC must also know the policies and procedures within the food establishment. If a PIC is not able to demonstrate knowledge, or is not onsite, the facility will receive a critical violation during an inspection. Please refer to chapter two of the Food Sanitation Rules to learn about the requirements of the PIC.



Food Service Inspection Process

All establishments must follow standard food-safety practices critical to the safety and quality of the food served. During health department inspections, the inspector compares the Food Sanitation Rules to the procedures at your establishment. The PIC can solicit help from the health department on how to prepare for an official regulatory inspection. The PIC can learn to identify every food safety problem, and how to correct it.

Employee Illness

Employee Illness

Infected food workers can spread a wide range of illnesses to others through food and utensils. Employees sick with vomiting or diarrhea must not work until symptoms have been gone for at least 24 hours.

Employees are required to report to the PIC when ill with any of the diseases listed below, or they live in the same household as a person with one of these diseases. An employee that has diarrhea, vomiting, jaundice, and sore throat with fever must also report this information to the PIC. The PIC is required to inform employees of this responsibility.

Most Common Illnesses from Employees

Foodborne Illnesses*	Common Symptoms				
	D	F	V	J	S
1. Hepatitis A virus		F		J	
2. Salmonella Typhi		F			
3. Shigella	D	F	V		
4. E. coli O157:H7	D				
5. Norwalk or Noro viruses	D	F	V		
6. Staphylococcus aureus	D		V		
7. Streptococcus pyogenes		F			S

Key: **D**=Diarrhea, **V**=Vomiting, **J**=Jaundice, **F**=Fever, **S**=Sore throat with Fever

Note: *The PIC is required to notify the county health department when an employee has Norovirus, Hepatitis A, Salmonella Typhi, Shigella or E. coli. O157:H7

Hand-Washing

Keeping Hands Clean

Food employees with dirty hands and/or dirty fingernails may contaminate the food being prepared. Any activity that may have contaminated the hands must be followed by thorough hand-washing as described in this manual (see pages 3 & 11).



Cleaning Procedure

Many employees fail to wash their hands as often as necessary and even those who do may use a poor technique. It takes vigorous rubbing with soap and running water for about 20 seconds to loosen soil and illness-causing organisms (germs).

Many diseases transferred through food may be harbored in the employee's intestinal tract and shed in the feces. Proper hand-washing after a bowel movement establishes a barrier against the transfer of organisms (germs) present in the feces.

Hand Sanitizers

Sanitizer dips or hand sanitizers are **not** approved handwash techniques and are not acceptable substitutes to hand-washing.

Fingernails

The requirement for fingernails to be trimmed, filed, and maintained addresses the ability to clean beneath the fingernails. Failure to remove fecal material from beneath the fingernails after a bowel movement can be a major source of illness causing organisms. Ragged fingernails may harbor harmful organisms.

Jewelry

Items of jewelry such as rings, bracelets, and watches may act as a hiding place for foodborne illness causing organisms (germs). An additional hazard associated with jewelry is the possibility that pieces of the item or the whole item itself may fall into the food being prepared. Hard foreign objects in food may cause medical problems for consumers, such as chipped and or broken teeth and internal cuts and lesions.

Cooking Temperatures

Cooking Temperatures

Different raw animal foods have to reach different temperatures to be done or safe. Use a metal-stem probe food thermometer to check temperatures while cooking to make sure that it gets done all the way inside.

Why Use a Food Thermometer?

A metal-stem probe food thermometer is the only reliable way to know that the temperature of food is hot enough to kill harmful bacteria. It also helps you to avoid overcooking. Each kitchen must have at least one accurate food thermometer that has a small enough probe to fit into thin food such as a thin meat patty.

Cooking Requirements for Specific Foods

The next page shows the temperature that specific foods must reach to be safe.



Cooking Requirements for Specific Foods

Animal Product	Minimum Temperature	What to Know?
Poultry, ground poultry	165°F (74°C) for 15 seconds	Stuffing should be cooked outside of poultry.
Stuffing, stuffed meats, casseroles and dishes combining raw and cooked food	165°F (74°C) for 15 seconds	Stuffing acts as an insulator, preventing heat from reaching the meat's center. Stuffing should be cooked separately.
Ground or flaked meats hamburger, ground pork, flaked fish, ground game animals, sausage, injected and pinned meats	155°F (68°C) for 15 seconds	Grinding meat mixes the organisms from the surface into the meat. Alternative minimum internal temperatures for ground meats: 150°F (66°C) for 1 minute 145°F (63°C) for 3 minutes
Pork, beef steaks, veal lamb, commercially raised game animals	145°F (63°C) for 15 seconds	This temperature is high enough to destroy Trichinella larvae that may have infested pork.
Beef or pork roasts	145°F (63°C) for 3 minutes	Alternative minimum internal cooking temperatures for beef and pork roasts: 130°F (54°C) for 121 minutes 134°F (57°C) for 47 minutes 138°F (59°C) for 19 minutes 140°F (60°C) for 12 minutes 142°F (61°C) for 8 minutes 144°F (62°C) for 5 minutes
Fish, foods containing fish, and seafood	145°F (63°C) for 15 seconds	Stuffed fish should be cooked to 165°F (74°C) for 15 seconds Fish that has been ground, chopped, or minced should be cooked to 155°F (68°C) for 15 seconds
Shell eggs for immediate service	145°F (63°C) for 15 seconds	Only take out as many eggs as you need. Never stack egg flats near the grill or stove. Eggs cooked for later service must be cooked to 155°F for 15 seconds and held at 135°F.
Foods cooked in microwave - meat, poultry, fish, eggs	165°F (74°C) let it stand for 2 minutes after cooking	Cover food, rotate or stir it halfway through the cooking process.

“Is It Done Yet?”




How to Use a Food Thermometer

- A thermometer that works best has a range of 0°F (-18°C) to 220°F (104°C).
- Use a thermometer with a smaller diameter probe on thin foods such as thin hamburger patties.
- Check the internal temperature of the food toward the end of the cooking time.
- Place the thermometer in the thickest part of the meat or in the center of the food to get a true reading. (Do not touch the bone with the stem of the thermometer to prevent a false reading).
- When taking temperatures of a large amount of food like a big piece of meat, be sure to take the temperature in two or more locations.
- Compare your thermometer reading to the Required Cooking Temperatures on the previous page to determine if your food has reached a safe temperature.
- Wash and sanitize the thermometer each time you check the temperature of a food.

Refrigerator Thermometer

Every refrigerator is required to have a thermometer. This thermometer must be located where it is easy to see when you open the refrigerator door. Every refrigerator should be operating at 41°F or less as indicated by the thermometer. If the thermometer reads above 41°F, then use a metal-stem probe food thermometer to check the temperature of food inside of the refrigerator with a food thermometer.




Types of Food Thermometers		
Name	Speed	Placement
Thermocouple (Most models can be calibrated)	2-5 seconds	1/4" or deeper in the food as needed
		
Thermistor (Some models can be calibrated)	10 seconds	At least 1/2" deep in the food
		
Instant-Read Bimetal (Most models can be calibrated)	15-20 seconds	2- 2 1/2" deep
		

Calibrating a Dial Food Thermometer

When you use a dial food thermometer you need to make sure the temperature it gives you is accurate. An easy way to do this is to use ice and water.

Pack a large cup to the top with crushed ice and water. Put the thermometer at least 2 inches into the ice slurry. After 30 seconds, read the dial. It should read 32°F (0°C). If it does not, you need to:

1. Leave it in the ice slurry. Add ice as it melts.
2. Use pliers or a wrench and turn the nut on the back of the thermometer until the needle reads 32°F (0°C).
3. Wait 30 seconds. Keep repeating these steps until the thermometer reads 32°F (0°C).





Cooling Hot Foods

With cooling of foods it is important to move the food temperature through the “Danger Zone” as quickly as possible to keep food safe.

Fresh is Best

You always take a chance that bacteria can grow and produce toxins when you cool food. It is safest to make food fresh each day, just before you serve it.

Speed is Important with Cooling

If you must make food in advance or save leftover food, cool it as **fast** as you can to prevent bacteria growth and toxin production. **Reheating will not destroy toxins.**

Cooling Solid Foods

When cooling solid cooked foods such as roast, turkey, and solid cuts of meat, be sure to:

1. Cut large roasts and turkeys into smaller portions. This will help them to cool faster.
2. Put all meats and other hot food in the refrigerator uncovered.

Cooling Soft/Thick Foods

Examples of soft/thick foods are refried beans, rice, potatoes, stews, chili, thick soup or thick sauces.

You can cool soft/thick foods by pouring food into a shallow metal pan. Use a sheet pan for very thick foods like refried beans.

Cooling thick food is not easy. Whenever possible, use a flat sheet pan and spread the food out as shallow as you can to speed up the cooling.

When cooling food in shallow metal pans, be sure to:

1. Pour hot food into shallow metal pans. The shallower the pan the faster the food will cool.
2. Stirring food speeds up cooling time.
3. Once food cools to 41°F (5°C), you can place food in a larger container and cover it.

Air Movement


Air in the refrigerator must be able to move around the food. The pans and dishes need to have space between them; do not crowd them. Do not stack pans on each other. Do not cover the food while it is cooling. A cover may be put on after the food has fully cooled.



Cooling Liquid Foods

You can use shallow metal pans or you can use the ice and water bath to cool thin soup and sauces. When cooling food with an ice bath, be sure to:

1. Close the drain in a large sink. Place the metal pot or pan of hot food in the sink. The sink drain must be indirectly plumbed.
2. Fill the sink with ice and cold water **up to the level of food in the pot or pan.**
3. Stir the soup or sauce often so that it cools all the way to the center. Ice paddles or cooling wands can be used to speed up the cooling process.
4. Add more ice as ice melts.
5. The food must reach 41°F (5°C).



Remember

You can choose several ways to cool food. No matter how you cool the food, it must drop from:

2 Hours	135°F (57°C) to 70°F (21°C) within two hours and then the temperature must drop from
4 Hours	70°F (21°C) to 41°F (5°C) within four hours.

Use a food thermometer to check the temperature while it is cooling. If it isn't cooling fast enough, you will need to do something else to speed up cooling.

Date Marking

Ready-to-eat potentially hazardous foods must be date marked with either the preparation date, use-by date, or date the commercial package was opened.

Seven Days

Ready-to-eat potentially hazardous foods must be date marked with either the preparation date, use-by date, or date the commercial package was opened.

The food can be stored for seven days when the refrigerator maintains 41°F (5°C) or colder. Food older than seven days must be discarded.

One Day

Food used within one day is not required to be date marked.

Pests

Cockroaches, flies, mice and rats can carry disease and cause damage. Prevention and control of these pests is essential.

Keep the inside and outside areas clean. Outside garbage must be contained in watertight containers with lids remaining closed when not in use. Exclude flies, especially during the warmer months, by screening open doors and windows screened with 1/16th of an inch mesh.

Pests can come into the facility through small holes or gaps under the door to the outside. A mouse can slip through a space of 1/4 inch. Block their entry by eliminating small holes and gaps under and around the door.

If you find pests inside your facility, contact a licensed pest control service.



Acknowledgements

Our Thanks to the California and Oregon Health Authorities,
Office of Environmental Public Health Foodborne Illness
Prevention Program for these materials. The eFoodcard Certificate
Program is FDA 2011 Code Compliant.